arXiv:1410.5700v1 [hep-ex] 21 Oct 2014

# Results for $\phi_2$ from Belle

*Pit Vanhoefer*
*Max-Plank-Institut für Physik*
*Föringer Ring 6*
*München 80805 GERMANY*

We present a summary of measurements sensitive to the CKM angle $\phi_2$, performed by the Belle experiment using the final data sample of $772\times 10^6$ $B\overline{B}$ pairs produced at the $\Upsilon(4S)$ resonance at the KEK asymmetric $e^+e^-$ collider. We discuss $CP$ asymmetries from the decay $B \to \pi^+\pi^-$ and briefly mention a preliminary measurement of the branching fraction of $B \to \pi^0\pi^0$ decays. Furthermore the measurement of the branching fraction of $B^0 \to \rho^0\rho^0$ decays and fraction of longitudinal polarization in this decay is presented. We use the results to constrain $\phi_2$ with isospin analyses in the $B \to \pi\pi$ and $B \to \rho\rho$ systems.

PRESENTED AT

CKM2014, the 8th International Workshop on the CKM Unitarity Triangle
Vienna, Austria

# 1 Introduction

One major precision test of the Standard Model (SM) is to validate the Cabibbo-Kobayashi-Maskawa (CKM) mechanism for violation of the combined charge-parity ($CP$) symmetry [1, 2]. This is one of the main purposes of the Belle experiment at KEK which has significantly contributed to proving the CKM scheme and constraining the unitarity triangle for $B$ decays to its current precision. Any deviation from unitarity would be a clear hint for physics beyond the SM. These proceedings give a summary of the experimental status of measurements of the CKM angle $\phi_2$ defined from the CKM matrix elements as $\phi_2 \equiv \arg(-V_{td}V_{tb}^*)/(V_{ud}V_{ub}^*)$.

The CKM angles can be determined by measuring the time-dependent asymmetry between $B^0$ and $\overline{B}^0$ decays into a common $CP$ eigenstate [3]. In the decay sequence, $\Upsilon(4S) \to B_{CP}B_{\rm Tag} \to f_{CP}f_{\rm Tag}$, where one of the $B$ mesons decays into a $CP$ eigenstate $f_{CP}$ at a time $t_{CP}$ and the other decays into a flavour specific final state $f_{\rm Tag}$ at a time $t_{\rm Tag}$, the time-dependent decay rate is given by

$$P(\Delta t, q) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}}\Big[1 + q(A_{CP}\cos\Delta m_d\Delta t + S_{CP}\sin\Delta m_d\Delta t)\Big], \qquad (1)$$

where $\Delta t \equiv t_{CP} - t_{\rm Tag}$ is the lifetime difference between the two $B$ mesons, $\Delta m_d$ is the mass difference between the mass eigenstates $B_H$ and $B_L$ and $q = +1(-1)$ for $B_{\rm Tag} = B^0(\overline{B}^0)$. The $CP$ asymmetry is given by

$$\frac{N(\overline{B} \to f_{CP}) - N(B \to f_{CP})}{N(\overline{B} \to f_{CP}) + N(B \to f_{CP})}, \qquad (2)$$

where $N(B(\overline{B}) \to f_{CP})$ is the number of events of a $B(\overline{B})$ decaying to $f_{CP}$, the asymmetry can be time-dependent. The parameters $A_{CP}$ and $S_{CP}$ describe direct and mixing-induced $CP$ violation, respectively *. All measurements presented here are based on Belle's final data set of $772 \times 10^6$ $B\overline{B}$ pairs.

# 2 The CKM Angle $\phi_2$

Decays proceeding via $b \to u\overline{u}d$ quark transitions such as $B^0 \to \pi\pi$, $\rho\pi$, $\rho\rho$ and $a_1(1260)\pi$, are directly sensitive to $\phi_2$. At tree level we expect $A_{CP} = 0$ and $S_{CP} = \sin 2\phi_2$. Possible penguin contributions can give rise of direct $CP$ violation, $A_{CP} \neq 0$ and also pollute the measurement of $\phi_2$, $S_{CP} = \sqrt{1 - A_{CP}^2}\sin(2\phi_2^{eff})$ where the observed $\phi_2^{eff} \equiv \phi_2 - \Delta\phi_2$ is shifted by $\Delta\phi_2$ due to different weak and strong phases from additional non-leading contributions. Despite this, it is possible to determine

*There exists an alternate notation where $C_{CP} = -A_{CP}$.




Figure 1: a) leading order and b) penguin feynman diagrams for $b \to u\overline{u}d$ transitions.

$\Delta\phi_2$ in $B^0 \to h^+h^-$ with an $SU(2)$ isospin analysis by considering the set of three $B \to hh$ decays where the $hh$s are either two pions or two longitudinally polarized $\rho s$, related via isospin symmetry [4]. The $B \to hh$ amplitudes obey the triangle relations,

$$A_{+0} = \frac{1}{\sqrt{2}}A_{+-} + A_{00}, \quad \overline{A}_{-0} = \frac{1}{\sqrt{2}}\overline{A}_{+-} + \overline{A}_{00}. \tag{3}$$

Isospin arguments demonstrate that $B^+ \to h^+h^0$ is a pure first-order mode in the limit of neglecting electroweak penguins, thus these triangles share the same base, $A_{+0} = \overline{A}_{-0}$, see Fig. 2 for an illustration. $\Delta\phi_2$ can then be determined from the difference between the two triangles. This method has an inherent 8-fold discrete ambiguity in the determination of $\phi_2$.




Figure 2: A sketch of the isospin triangle.

## 3 The Decay $B^0 \to \pi^+\pi^-$

A seven-dimensional extended maximum likelihood fit to the difference of the energy of the reconstructed $B$ meson to the beam-energy ($\Delta E$), the beam-energy constraint

$B$ mass ($M_{\rm bc}$), two likelihoods for each charged track to be a Kaon assuming a $\pi^{\pm}$ mass hypothesis ($L^{\pm}_{K/pi}$), a event shape-dependent fisher discriminant ($\mathcal{F}_{B\overline{B}/q\overline{q}}$) which separates $B\overline{B}$ (spherical) from $q\overline{q}$ (two-jet like) events, the life-time difference of $B_{CP}$ and $B_{\rm tag}$ ($\Delta t$) and the flavor of $B_{\rm tag}$ ($q$), with $q = +1(-1)$ for $B_{\rm tag} = B^0(\overline{B^0})$ was performed to simultaniously obtain the yields and $CP$ asymmetries of $B \to h^+h^-$ ($h = K, \pi$) decays [5]. The measurement of the $CP$ parameters in $B^0 \to \pi^+\pi^-$ decays yield $\mathcal{S}_{CP} = -0.636 \pm 0.082$ (stat) $\pm 0.027$ (syst) and $\mathcal{A}_{CP} = 0.328 \pm 0.061$ (stat) $\pm$ 0.027 (syst). The $M_{\rm bc}$ and $\Delta t$ distributions are shown in Fig. 3. Belle excludes the range $\phi_2 \notin [23.8°, 66.8°]$ at the $1\sigma$ level by performing an isospin analysis to remove the penguin contribution, see Fig. 4 a). The amount of direct $CP$ violation was found to be smaller compared to the previous measurement at Belle [6]. The previous result was confirmed with the previous data set of $535 \times 10^6$ $B\overline{B}$ pairs. While their was already good agreement in previous measurements of $\mathcal{S}_{CP}$, the drop in $\mathcal{A}_{CP}$ leads to a in better agreement with other experiments [7, 8]. This is currently the most precise measurement of $CP$ violation in $B^0 \to \pi^+\pi^-$ decays.




Figure 3: a) $M_{\rm bc}$ distribution and b) $\Delta t$ distribution for each flavour tag and the fit result on top. b) also shows the resulting $CP$ asymmetry for $B^0 \to \pi^+\pi^-$. Mixing-induced $CP$ violation can be clearly seen in the asymmetry plots and the height difference in the $\Delta t$ projection indicates direct $CP$ violation.

# 4 The Decay $B^0 \to \pi^0\pi^0$

We only report on this mode briefly as it is discussed in detail in another presentation (M.Seviour, WG $V$). This decay is reconstructed from four $\gamma$s making this measure-




Figure 4: Scans of $\phi_2$ from an isospin analysis in the a) $B \to \pi\pi$ and b) $B \to \rho\rho$ system, the dashed line corresponds to the one $\sigma$ level.

ment experimentally quite challenging. A fit to $\Delta E$, $M_{\rm bc}$ and $\mathcal{F}_{B\overline{B}/q\overline{q}}$ is performed and a preliminary branching fraction of $\mathcal{B}(B \to \pi^0\pi^0) = (0.9 \pm 0.12\ \text{stat} \pm 0.10\ \text{syst})$ is obtained. It is planned to supersede this measurement with one including a measurement of direct $CP$ violation in this mode. For next generation $B$ factories photon conversion makes vertexing possible and allows to perform a measurement of the time-dependent $CP$ asymetries.

# 5 The Decay $B^0 \to \rho^0\rho^0$

Having a decay into two vector particles, an angular analysis has to be performed in order to separate the pure $CP$-even states with longitudinal polarization from the transverse polarized mixture of $CP$-even and odd states for the isopsin analysis. The presence of multiple, largely unknown backgrounds with the same four charged pions final state as $B^0 \to \rho^0\rho^0$ make this decay quite difficult to isolate and interference between the various $4\pi$ modes needs to be considered. Similar as for $B^0 \to \rho^+\rho^-$ decays [9, 10, 11], the decay $B \to \rho\rho$ is naively expected to be polarized dominantly longitudinally. However, color-suppressed $B$ decays into light vectors are especially difficult to predict as the transverse polarized amplitude is not calculable [12] . Besides updating to the full data set, two helicity angles $\cos\Theta_H^{1,2}$, one for each $\rho^0$, are added to the fit. The angles, defined in the helicity basis, are powerful in separating the different backgrounds and allow one to measure the fraction $f_L$ of longitudinal polarization. Belle obtains $\mathcal{B}(B^0 \to \rho^0\rho^0) = (1.02 \pm 0.30\ (\text{stat}) \pm 0.15\ (\text{syst})) \times 10^{-6}$ with a fraction of longitudinal polarization, $f_L = 0.21^{+0.18}_{-0.22}\ (\text{stat}) \pm 0.15\ (\text{syst})$ [13]. Signal enhanced distributions onto $\Delta E$, $m(\pi^+\pi^-)$ and one of the helicty angles, each with the fit result on top, are shown in Fig. 5. Since this mode is found to decay dominantly into trans-

versely polarized $\rho^0$s, a measurement of the $CP$ asymmetries has not been performed. However, the size of the amplitude of the decays into longitudinally polarized $\rho^0$s from this measurement has been used in an isospin analysis together with other Belle measurements [10, 11, 14] (longitudinal polarization only). The resulting constraint most consistent with other measurements of the CKM triangle is $\phi_2 = (84.9 \pm 13.5)^\circ$. The relatively small amplitude of $B^0 \to \rho^0\rho^0$ makes the isospin analysis in the $B \to \rho\rho$ less ambiguous. Fig. 4 b) shows $\phi_2$ scan from the isospin analysis. Comparing these results with the ones obtained by BaBar, we find good agreement in the branching fraction of $B^0 \to \rho^0\rho^0$ decays, while there is a $2.1\sigma$ discrepancy in the fraction of longitudinal polarization. BaBar finds $f_L = 0.75^{+0.11}_{-0.14}$ (stat) $\pm 0.04$ (syst) [15]. This measurement is still statistically (and systematically) limited, hence further studies with more statistics would be very interesting and hopefully will solve this tensions.

# 6 Summary

We have presented recent measurements from Belle sensitive to the CKM phase $\phi_2$ using the full data set. Measurements of the $CP$ asymmetries in $B \to \pi^+\pi^-$ and the branching fraction of $B \to \rho^0\rho^0$, together with fraction of longitudinal polarized $\rho^0$s in this decay were presented. The data are used to constrain $\phi_2$ with an $SU(2)$ isospin analysis for each isospin triplet. The current world averages of $\phi_2$ as computed by the CKMfitter [16] (including the results presented) and UTfit [17] collaborations are $\phi_2 = (88.5^{+4.7}_{-4.4})^\circ$ and $\phi_2 = (89.1 \pm 3.0)^\circ$, respectively. Furthermore we presented a preliminary update of the branching fraction of $B^0 \to \pi^0\pi^0$ decays. With BelleII being built and LHCb operating, the next generation of $B$ physics experiments are expected to further reduce the uncertainty of the CKM observables, e.g. the uncertainty of $\phi_2$ is expected to be reduced to $\sim 1^\circ$ [18].

# References


[1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).

[2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

[3] I. Bigi and A. Sanda, *CP Violation*, Cambridge University Press, Cambridge (2009).

[4] M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990).

[5] J.Dalseno *et al.* (Belle Collab.), Phys. Rev. D **88**, 092003 (2013).

[6] H. Ishino *et al.* (Belle Collab.), Phys. Rev. Lett. **98**, 211801 (2007).

[7] J. P. Lees *et al.* (BaBar Collab.), arXiv:1206.3525v1 [hep-ex].

[8] ,R. Aaij *et al.* (LHCb Collab.) JHEP 1310 **183** (2013).

[9] B. Aubert *et al.* (BaBar Collab.), Phys. Rev. D **76**, 052007 (2007).

[10] A. Somov *et al.* (Belle Collab.), Phys. Rev. Lett. **96**, 171801 (2006).

[11] A. Somov *et al.* (Belle Collab.), Phys. Rev. D **76**, 011104 (2007).

[12] M. Beneke, J. Rohrer and D. Yang, Nucl. Phys. B **774**, 64-101 (2007).

[13] P.Vanhoefer *et al.* (Belle Collab.), Phys. Rev. D **89**, 072008 (2014).

[14] J. Zhang *et al.* (Belle Collab.), Phys. Rev. L **91**, 221801 (2006).

[15] B. Aubert *et al.* (BaBar Collab.), Phys. Rev. D **78**, 071104 (2008).

[16] http://ckmfitter.in2p3.fr

[17] http://www.utfit.org

[18] T. Aushev *et al.*, arXiv:1002.5012 (2010).




Figure 5: Signal enhanced distributions of a) $\Delta E$, b) $m(\pi^+\pi^-)$ and c,d) $\cos\Theta_H$ with the fit result on top. d) shows a distribution where the contributions from the remaining four-pion final states are reduced by a tight cut on the $m(\pi^+\pi^-)$-$m(\pi^+\pi^-)$ distribution, demonstrating the necessity of a transversaly polarized component. The shaded red area and the long dashed orange histogram are the $B^0 \to \rho^0\rho^0$ and $f_0\rho^0$ contributions, respectively. Furthermore, all four pion final states are shown in dashed cyan, the entire ($B\overline{B}$) background in dashed green (dark green) and the full PDF in blue.